

The Journal of Physiology

Limitations to systemic and locomotor limb muscle oxygen delivery and uptake during maximal exercise in humans

Stefan P Mortensen, Ellen A Dawson, Chie C Yoshiga, Mads K Dalsgaard, Rasmus Damsgaard, Niels H Secher and José González-Alonso

J. Physiol. 2005;566;273-285; originally published online Apr 28, 2005;

DOI: 10.1113/jphysiol.2005.086025

This information is current as of October 11, 2007

This is the final published version of this article; it is available at:

<http://jp.physoc.org/cgi/content/full/566/1/273>

This version of the article may not be posted on a public website for 12 months after publication unless article is open access.

The Journal of Physiology Online is the official journal of The Physiological Society. It has been published continuously since 1878. To subscribe to *The Journal of Physiology Online* go to: <http://jp.physoc.org/subscriptions/>. *The Journal of Physiology Online* articles are free 12 months after publication. No part of this article may be reproduced without the permission of Blackwell Publishing: JournalsRights@oxon.blackwellpublishing.com

Limitations to systemic and locomotor limb muscle oxygen delivery and uptake during maximal exercise in humans

Stefan P. Mortensen¹, Ellen A. Dawson^{1,2}, Chie C. Yoshiga^{1,2}, Mads K. Dalsgaard^{1,2}, Rasmus Damsgaard¹, Niels H. Secher^{1,2} and José González-Alonso¹

¹The Copenhagen Muscle Research Centre and ²Department of Anaesthesia, Rigshospitalet, University of Copenhagen, Denmark

Reductions in systemic and locomotor limb muscle blood flow and O₂ delivery limit aerobic capacity in humans. To examine whether O₂ delivery limits both aerobic power and capacity, we first measured systemic haemodynamics, O₂ transport and O₂ uptake (\dot{V}_{O_2}) during incremental and constant (372 ± 11 W; 85% of peak power; mean \pm S.E.M.) cycling exercise to exhaustion ($n = 8$) and then measured systemic and leg haemodynamics and \dot{V}_{O_2} during incremental cycling and knee-extensor exercise in male subjects ($n = 10$). During incremental cycling, cardiac output (\dot{Q}) and systemic O₂ delivery increased linearly to 80% of peak power ($r^2 = 0.998$, $P < 0.001$) and then plateaued in parallel to a decline in stroke volume (SV) and an increase in central venous and mean arterial pressures ($P < 0.05$). In contrast, heart rate and \dot{V}_{O_2} increased linearly until exhaustion ($r^2 = 0.993$; $P < 0.001$) accompanying a rise in systemic O₂ extraction to $84 \pm 2\%$. In the exercising legs, blood flow and O₂ delivery levelled off at 73–88% of peak power, blunting leg \dot{V}_{O_2} per unit of work despite increasing O₂ extraction. When blood flow increased linearly during one-legged knee-extensor exercise, \dot{V}_{O_2} per unit of work was unaltered on fatigue. During constant cycling, \dot{Q} , SV, systemic O₂ delivery and \dot{V}_{O_2} reached maximal values within ~ 5 min, but dropped before exhaustion ($P < 0.05$) despite increasing or stable central venous and mean arterial pressures. In both types of maximal cycling, the impaired systemic O₂ delivery was due to the decline or plateau in \dot{Q} because arterial O₂ content continued to increase. These results indicate that an inability of the circulatory system to sustain a linear increase in O₂ delivery to the locomotor muscles restrains aerobic power. The similar impairment in SV and O₂ delivery during incremental and constant load cycling provides evidence for a central limitation to aerobic power and capacity in humans.

(Resubmitted 9 March 2005; accepted 27 April 2005; first published online 28 April 2005)

Corresponding author J. González-Alonso: The Copenhagen Muscle Research Centre, Rigshospitalet, Section 7652, Blegdamsvej 9, DK-2100 Copenhagen Ø, Denmark. Email: jga@cmrc.dk

From rest to maximal exercise, the cardiovascular system must adjust to the increasing metabolic demand by ensuring the delivery of O₂ and substrates to all body cells, in particular to the myocytes, without compromising arterial pressure. The prevailing theory is that the cardiovascular system responds to exercise of increasing intensity up to aerobic power (maximal oxygen uptake ($\dot{V}_{O_{2,max}}$)) by increasing systemic and contracting skeletal muscle O₂ delivery in proportion to the rise in O₂ demand, while increasing the perfusion pressure (Holmgren, 1956; Åstrand *et al.* 1964; Ekelund & Holmgren, 1967; Higginbotham *et al.* 1986; Reeves *et al.* 1990; Pawelczyk *et al.* 1992; Rowell, 1993; Rowell *et al.* 1996). However, recent findings during constant load maximal cycling

exercise indicate that, following early adjustments that allow $\dot{V}_{O_{2,max}}$ to be reached and maintained for 1–2 min, cardiac output (\dot{Q}), leg blood flow (LBF), perfusion pressure and O₂ delivery decline leading to reductions in $\dot{V}_{O_{2,max}}$ and leg \dot{V}_{O_2} despite increases in O₂ extraction (González-Alonso & Calbet, 2003; González-Alonso *et al.* 2004). These findings suggest that a reduction in systemic and locomotive skeletal muscle O₂ delivery limits aerobic and maximal endurance capacity in trained individuals. The question then arises whether systemic and locomotor limb muscle O₂ delivery is also impaired during incremental exercise and whether this could explain the plateau or decline in \dot{V}_{O_2} observed sometimes before exhaustion (Åstrand, 1952; Taylor *et al.*

1955; Mitchell *et al.* 1958; Froelicher *et al.* 1972; Pollock *et al.* 1976; Meyers *et al.* 1990; Knight *et al.* 1992; Day *et al.* 2003).

Indirect evidence indicates that O₂ delivery might impose a limitation to \dot{V}_{O_2} at intensities close to $\dot{V}_{O_{2,max}}$. Firstly, in humans the increase in \dot{V}_{O_2} per unit of work is attenuated at high compared to low exercise intensities (Hill & Lupton, 1923; Åstrand & Saltin, 1961; Whipp & Wasserman, 1972). Secondly, the increase in \dot{Q} per litre increase in \dot{V}_{O_2} is attenuated at intensities above 40–70% of $\dot{V}_{O_{2,max}}$ in humans, as in miniature swine (Saltin, 1964; Åstrand *et al.* 1964; Armstrong *et al.* 1987). These early observations lack conclusive support and are therefore not widely accepted. The general belief is that \dot{Q} and O₂ delivery increase linearly from rest to $\dot{V}_{O_{2,max}}$, implying that O₂ delivery to locomotor limb muscle does not limit \dot{V}_{O_2} (Asmussen & Nielsen, 1952; Chapman *et al.* 1960; Bevegård *et al.* 1963; Rowell *et al.* 1964; Grimsby *et al.* 1966; Poliner *et al.* 1980; Higginbotham *et al.* 1986). This theory, however, is based on linear regression analysis of \dot{Q} data that have not been normalized and on the assumption that systemic haemodynamics closely reflect skeletal muscle haemodynamics. At the skeletal muscle level, not only restrictions in the systemic supply of O₂, but also limitations in diffusive O₂ transport from the muscle capillary to the mitochondrial cytochrome and oxidative capacity of mitochondria could restrict $\dot{V}_{O_{2,max}}$ (Roca *et al.* 1989). In favour of a dominant O₂ supply limitation, quadriceps muscle blood flow and \dot{V}_{O_2} might reach ~ 2.3 and ~ 0.35 l kg⁻¹ min⁻¹ during maximal knee-extensor exercise, suggesting that only 10–15 kg of muscle needs to be recruited during whole body exercise to surpass the capacity of the human circulation to deliver O₂ (Andersen & Saltin, 1985). Whether the O₂ delivery to muscles and uptake of the quadriceps femoris are restricted during whole body compared to knee-extensor exercise remains unknown.




Figure 1. Core temperature during the constant and incremental protocols

Oesophageal temperature at rest, during submaximal and maximal exercise, and during 10 min of recovery, in incremental (●) and constant (○) exercise protocols. Data are means \pm S.E.M. for 8 subjects.

To investigate the contribution of the O₂ transport system to $\dot{V}_{O_{2,max}}$, we determined: (1) whether systemic O₂ delivery imposes a limitation to aerobic power and capacity, (2) whether locomotor limb blood flow and O₂ delivery are impaired during incremental exercise to exhaustion to the extent that they compromise limb \dot{V}_{O_2} , and (3) whether quadriceps muscle blood flow and \dot{V}_{O_2} are lower during maximal exercise with a large compared to a small muscle mass. To accomplish these aims, we first measured systemic haemodynamics, O₂ transport and \dot{V}_{O_2} during incremental and constant cycling exercise to exhaustion in trained male subjects and then measured systemic and exercising leg haemodynamics, O₂ transport and \dot{V}_{O_2} during incremental cycle and knee-extensor exercise to exhaustion in another group of active male subjects. We hypothesized that restrictions in O₂ supply to locomotor limb muscles imposes a limitation to aerobic power and capacity in humans.

Methods

Eighteen endurance-trained or recreationally active male subjects participated in two studies. They had a mean (\pm s.d.) age of 27 ± 3 years, body weight of 81.3 ± 9.6 kg, height of 185 ± 9 cm, maximal heart rate (HR) of 192 ± 7 beats min⁻¹ and $\dot{V}_{O_{2,max}}$ of 4.80 ± 0.46 l min⁻¹. The subjects were informed of any risks and discomforts associated with the experiments before giving written consent to participate. The study was approved by the Ethics Committee of Copenhagen (KF 01-230/00) and conducted in accordance with the guidelines of the *Declaration of Helsinki*.

In the first study and on the first visit to the laboratory, eight endurance-trained subjects performed incremental exercise on a cycle ergometer (Excalibur, Lode, The Netherlands) to determine $\dot{V}_{O_{2,max}}$, maximal HR and peak power. Thereafter, they completed four high-intensity training sessions on the cycle ergometer. During the last session, they carried out the same protocol as during the main experiment involving incremental (INC) and constant (CON) maximal exercise separated by one hour of recovery, while continuous measures of \dot{V}_{O_2} , HR and oesophageal temperature were obtained. For the invasive experiment, the subjects arrived at the laboratory one hour prior to the experiment after a light breakfast. Catheters were placed into the brachial artery and an antecubital vein, with the latter catheter being advanced to the right atrium. Following 30 min of supine rest, the subjects completed INC and CON on the cycle ergometer preceded by a 15 min warm-up period (146 ± 6 W; $<50\%$ $\dot{V}_{O_{2,max}}$) and 3 min of rest, followed by 10 min of recovery. Throughout the protocol, the arms rested on aerobars simulating the position that cyclists adopt during a time-trial. During the resting and recovery periods, the subjects were allowed to move their legs (0 W). During INC, the workload was

increased every minute using a computerized system to elicit 20, 40, 60, 80, 90, 95 and 100% of peak power. During CON, the intensity resulted in exhaustion within 5–7 min and $\dot{V}_{O_2\max}$ within 4–5 min (i.e. at 372 ± 11 W

or at 85% of the 438 ± 13 W peak power of the initial incremental test). The order of the two trials was randomly assigned and counterbalanced across the subjects. Exercise was performed under thermoneutral conditions ($\sim 20^\circ\text{C}$)




Figure 2. Central haemodynamics during the constant and incremental protocols

Cardiac output, heart rate, stroke volume, central venous and arterial pressure, systemic vascular conductance, arterial O_2 content, systemic O_2 delivery, a-v O_2 difference and \dot{V}_{O_2} at rest, during submaximal and maximal exercise, and during 10 min of recovery, in incremental (●) and constant load (○) exercise. Data are means \pm S.E.M. for 8 subjects, except arterial and central venous pressures and systemic vascular conductance for which data represent 7 subjects. * Lower than the value after 22 min when cycling at 80% of peak power, $P < 0.05$. ‡ Lower than the peak values observed after 20–24 min of constant load maximal cycling, $P < 0.05$.

with fans directed against the back and the side of the subjects.

Ten subjects participated in the second investigation which was aimed at determining whether convective O_2 transport and limb muscle \dot{V}_{O_2} are compromised during incremental cycle exercise to exhaustion and whether quadriceps muscle blood flow and \dot{V}_{O_2} are lower during maximal cycle compared to maximal knee-extensor

exercise. In these subjects, an additional catheter was inserted into the femoral vein 2 cm from the inguinal ligament to allow for blood sampling and measurements of LBF (Andersen & Saltin, 1985). During both types of incremental exercise, the workload was increased every 1.5 min to elicit 25, 50, 75, 90 and 100% of peak power.

In study 1, blood samples (1–5 ml) were drawn simultaneously from the brachial artery and the right




Figure 3. Central haemodynamics during incremental exercise to exhaustion

Cardiac output, heart rate, stroke volume, arterial (●) and central venous (▼) pressure, systemic vascular conductance, arterial O_2 content, systemic O_2 delivery, systemic a-v O_2 difference, systemic O_2 extraction and pulmonary \dot{V}_{O_2} during incremental exercise to exhaustion plotted against the relative increase in power output. Data are means \pm s.e.m. for 8 subjects. * Lower than 80% of peak power, $P < 0.05$.

atrium at rest in the supine and upright positions, during the warm-up (4, 10 and 15 min), immediately before the start of maximal exercise, during maximal exercise and during the recovery (1, 3, 6 and 10 min). In INC, blood samples were drawn after 45 s of each workload and at exhaustion. In CON, they were drawn after 0.75, 1.5, 3, 4 and 5 min and before exhaustion. In study 2, blood samples were drawn simultaneously from the brachial artery, right atrium and the femoral vein after 45 s at each workload, and LBF was measured after 1 min.

Throughout the studies, pulmonary \dot{V}_{O_2} was measured online (Medgraphics CPX/D, Saint Paul, MN, USA; study 1 and Cosmed Quark b², Italy; study 2). During the invasive experiments, HR was obtained from an electrocardiogram while arterial and central venous pressures were monitored with transducers positioned at heart level (Pressure Monitoring Kit, Baxter). The LBF was measured by the constant-infusion thermodilution method (Andersen & Saltin, 1985; González-Alonso *et al.* 2000b), while \dot{Q} was calculated using the Fick principle ($\dot{Q} = \dot{V}_{O_2}/\text{arteriovenous (a-v) } O_2 \text{ difference}$), assuming negligible differences in blood oxygenation between the right atrium and the pulmonary artery (Barratt-Boyes & Wood, 1956). The \dot{Q} data obtained in one subject using the direct Fick principle confirmed the \dot{Q} results. Stroke volume (SV) was the quotient between \dot{Q} and HR, and systemic and leg vascular conductance were the quotients between \dot{Q} and LBF and the perfusion pressure. Perfusion pressure was the difference between mean arterial (MAP) and central venous pressures and pulse pressure was that between the systolic and diastolic blood pressure. The left ventricular contractility index dP/dt_{\max} was calculated as the peak systolic value of the first derivative of the arterial pressure curve over 20 cardiac cycles. For systemic O_2 delivery, \dot{Q} was multiplied by the arterial O_2 content whereas systemic O_2 extraction was the ratio between the systemic a-v O_2 difference and the arterial O_2 content. Blood gases, haemoglobin, glucose and lactate concentrations were measured using an ABL700 analyser (Radiometer, Copenhagen, Denmark). Oesophageal temperature was measured with a thermocouple (MOV-A, Ellab, Copenhagen, Denmark) inserted through the nasal passage at a distance equal to one-fourth of the subject's standing height, and HR was measured with a Polar Sports Tester (Polar Electro). In study 1, blood gas variables were corrected for the temperature measured during the non-invasive trials, whereas in study 2 the correction was made from the femoral venous blood temperature. Leg muscle mass was calculated from the whole-body dual-energy X-ray absorptiometry scanning (Prodigy, General Electrics Medical Systems, WI, USA) as lean mass of the region. Quadriceps femoris muscle mass was calculated using the antropometric method, as described by Anderson & Saltin, 1985.

Statistical analysis

A one-way repeated measures analysis of variance (ANOVA) was performed to test significance within and between the two trials. Following a significant *F* test, pair-wise differences were identified using Tukey's honestly significant difference (HSD) *post hoc* procedure. To determine whether exhaustion during the constant maximal exercise was preceded by reductions in \dot{Q} , SV and O_2 delivery, final values were compared with peak values during exercise using one-way repeated measures ANOVA with Tukey's HSD *post hoc* procedure. The significance level was set at $P < 0.05$ and data are means \pm s.e.m. unless indicated otherwise.




Figure 4. Relationship between cardiac output and systemic \dot{V}_{O_2} during incremental exercise to exhaustion

Cardiac output, stroke volume and systemic O_2 delivery plotted against the increases in \dot{V}_{O_2} during incremental exercise. The plateau in cardiac output was due to a concomitant decline in stroke volume. Data are means \pm s.e.m. for 8 subjects. * Lower than the stroke volume value observed at a systematic \dot{V}_{O_2} of 3.9 L min⁻¹ when cycling at 80% of peak power, $P < 0.05$.

Table 1. Blood variables at rest, during incremental exercise to exhaustion, and after 10 min of recovery

	Rest	Incremental exercise (% peak power)							10 min recovery
		20%	40%	60%	80%	90%	95%	100%	
Haemoglobin (g l⁻¹)									
a	150 ± 4	153 ± 3	155 ± 3	156 ± 2*	157 ± 2*	158 ± 3*	160 ± 3*	162 ± 3*	156 ± 3*
v	151 ± 4	151 ± 4	152 ± 4	153 ± 4	158 ± 3*	159 ± 3*	161 ± 4*	163 ± 4*	154 ± 3
P_{O₂} (mmHg)									
a	120 ± 4	101 ± 2*	95 ± 2*	92 ± 4*	91 ± 3*	93 ± 3*	96 ± 3*	102 ± 3*	118 ± 2
v	40 ± 2	30 ± 2*	26 ± 1*	22 ± 1*	21 ± 1*	20 ± 1*	20 ± 1*	21 ± 1*	48 ± 2*
O₂ saturation (%)									
a	98.8 ± 0.1	98.0 ± 0.1	97.5 ± 0.2*	97.1 ± 0.3*	96.6 ± 0.3*	96.2 ± 0.3*	95.4 ± 0.5*	95.0 ± 0.5*	97.3 ± 0.2*
v	71.3 ± 2.3	50.4 ± 2.7*	37.6 ± 3.0*	30.4 ± 2.1*	25.6 ± 2.1*	20.9 ± 1.8*	17.2 ± 1.7*	14.8 ± 1.5*	69.8 ± 1.2
O₂ content (ml l⁻¹)									
a	202 ± 5	203 ± 4	205 ± 3	205 ± 3	206 ± 3	206 ± 3	208 ± 4	210 ± 4*	207 ± 4
v	146 ± 7	103 ± 7*	78 ± 7*	63 ± 5*	55 ± 5*	45 ± 4*	38 ± 4*	33 ± 4*	146 ± 5
P_{CO₂} (mmHg)									
a	41 ± 1	42 ± 1	42 ± 1	42 ± 0	42 ± 1	41 ± 1	39 ± 1	37 ± 1*	33 ± 1*
v	50 ± 1	53 ± 1	56 ± 1	61 ± 1*	70 ± 1*	77 ± 2*	83 ± 2*	87 ± 3*	43 ± 1*
pH									
a	7.40 ± 0.01	7.40 ± 0.00	7.39 ± 0.00	7.38 ± 0.01	7.36 ± 0.01*	7.32 ± 0.01*	7.26 ± 0.01*	7.20 ± 0.01*	7.19 ± 0.01*
v	7.35 ± 0.01	7.35 ± 0.00	7.33 ± 0.01	7.30 ± 0.01*	7.24 ± 0.01*	7.17 ± 0.01*	7.10 ± 0.01*	7.03 ± 0.01*	7.16 ± 0.01*
Lactate (mmol l⁻¹)									
a	1.0 ± 0.1	1.0 ± 0.1	1.3 ± 0.1	2.0 ± 0.2	4.2 ± 0.3*	7.7 ± 0.3*	11.9 ± 0.4*	16.4 ± 0.7*	14.3 ± 0.6*
v	1.0 ± 0.1	1.0 ± 0.1	1.7 ± 0.4	2.4 ± 0.4	4.3 ± 0.3*	8.0 ± 0.4*	12.1 ± 0.5*	16.8 ± 0.8*	13.7 ± 0.8*
Glucose (mmol l⁻¹)									
a	5.3 ± 0.3	5.3 ± 0.3	5.3 ± 0.3	5.4 ± 0.3	5.3 ± 0.3	5.2 ± 0.3	5.2 ± 0.3	5.0 ± 0.3	6.2 ± 0.2*
v	5.2 ± 0.3	5.2 ± 0.3	5.2 ± 0.3	5.2 ± 0.3	5.3 ± 0.3	5.2 ± 0.3	5.1 ± 0.3	4.9 ± 0.3	6.1 ± 0.2*

Values are means ± S.E.M. for 8 subjects. a, arterial. v, right atrium. * Different from rest, $P < 0.05$. P_{O_2} , P_{CO_2} and pH values were corrected for changes in blood temperature.

Results

Performance and maximal O₂ uptake

No significant differences in endurance, $\dot{V}_{O_{2max}}$ or peak power were observed between the non-invasive and the invasive experiments. In INC, time to fatigue was 6.45 ± 0.2 and 6.96 ± 0.1 min during the non-invasive and invasive experiments, respectively, accompanied by a similar $\dot{V}_{O_{2max}}$ (4.81 ± 0.12 and 4.75 ± 0.15 l min⁻¹, respectively) and peak power (440 ± 16 and 446 ± 13 W, respectively). Similarly, in CON, time to fatigue was 7.01 ± 0.23 and 6.87 ± 0.50 min during the non-invasive and invasive experiments, respectively, and $\dot{V}_{O_{2max}}$ was 4.82 ± 0.12 versus 4.75 ± 0.13 l min⁻¹. In INC the workload elicited 20 ± 0, 39 ± 0, 59 ± 1, 79 ± 1, 88 ± 15, 93 ± 1 and 100 ± 0% of peak power during the invasive experiment. Oesophageal temperature increased from $\sim 37.8^\circ\text{C}$ at the onset of exercise to 39.5 ± 0.1 and $39.9 \pm 0.1^\circ\text{C}$ in INC and CON, respectively (Fig. 1).

Systemic haemodynamics, O₂ transport and O₂ uptake

No differences in systemic haemodynamics, O₂ transport or \dot{V}_{O_2} were observed at rest or during the 15 min of

submaximal exercise (Fig. 2). In INC, \dot{Q} increased linearly to 80% of peak power ($r^2 = 0.998$; $P < 0.001$) and then plateaued (Figs 2–4). In CON, \dot{Q} increased during the first 1.5 min, reached a peak value of 27.1 ± 1.1 l min⁻¹ after 4.6 ± 0.6 min (range 3–6 min) and then declined 1.9 ± 0.5 l min⁻¹ before exhaustion ($P < 0.05$). The observed plateau in \dot{Q} above 80% of peak power in INC and the drop during CON were due to a fall in SV (20 ± 3 and 27 ± 6 ml beat⁻¹ in INC and CON, respectively), because HR continued to increase to exhaustion (190 ± 2 and 192 ± 2 beats min⁻¹, respectively). In both INC and CON, central venous pressure increased from -3 mmHg at rest to 2–5 mmHg at exhaustion. In INC, MAP increased from 94 ± 4 mmHg at the start of exercise to 136 ± 7 mmHg at exhaustion. In contrast, in CON, MAP stabilized at ~ 128 mmHg after 1.5 min. In INC, perfusion pressure increased from 98 ± 3 to 131 ± 6 mmHg at exhaustion accompanying an increase in pulse pressure from 59 ± 4 to 144 ± 4 mmHg. In contrast, perfusion pressure in CON increased from 100 ± 4 to 127 ± 6 mmHg after 3 min and remained stable thereafter. Similarly, pulse pressure increased from 60 ± 3 to 137 ± 4 mmHg after 3 min. In INC, systemic vascular conductance increased to $\sim 80\%$ of peak power (range 60–90%) and then declined ($P < 0.05$). In CON, vascular conductance

Table 2. Blood variables at rest, during constant load maximal exercise, and after 10 min of recovery

	Rest	Constant load exercise (min)					10 min recovery	
		0.8	1.5	3	4	5		6.9 ± 0.5
Haemoglobin (g l⁻¹)								
a	151 ± 3	155 ± 3	159 ± 3*	160 ± 4*	165 ± 3*	166 ± 3*	169 ± 3*	158 ± 4*
v	155 ± 3	158 ± 3	161 ± 4	163 ± 3	165 ± 3*	166 ± 3*	165 ± 3*	157 ± 5
P_{O₂} (mmHg)								
a	118 ± 4	91 ± 4*	91 ± 3*	92 ± 4*	94 ± 4*	96 ± 4*	101 ± 4*	117 ± 3
v	40 ± 1	22 ± 1*	20 ± 1*	20 ± 1*	20 ± 1*	20 ± 1*	20 ± 1*	49 ± 2*
O₂ saturation (%)								
a	98.8 ± 0.2	97.1 ± 0.5	96.6 ± 0.3*	96.1 ± 0.6*	94.7 ± 0.6*	94.3 ± 0.9*	93.0 ± 1.0*	96.9 ± 0.4*
v	70.8 ± 2.4	31.7 ± 2.8*	22.9 ± 2.3*	18.7 ± 1.7*	16.6 ± 1.6*	15.1 ± 1.8*	12.5 ± 1.2*	68.8 ± 1.3
O₂ content (ml l⁻¹)								
a	204 ± 4	205 ± 4	209 ± 5	209 ± 6	213 ± 4*	213 ± 4*	213 ± 4*	209 ± 5
v	149 ± 7	68 ± 7*	50 ± 6*	42 ± 4*	37 ± 4*	34 ± 4*	29 ± 3*	146 ± 5
P_{CO₂} (mmHg)								
a	41 ± 1	41 ± 1	41 ± 1	38 ± 1	37 ± 1	36 ± 1*	34 ± 1*	33 ± 1*
v	51 ± 1	58 ± 2*	71 ± 2*	76 ± 2*	78 ± 2*	79 ± 2*	82 ± 3*	43 ± 1*
pH								
a	7.41 ± 0.01	7.39 ± 0.01	7.35 ± 0.01	7.27 ± 0.01*	7.23 ± 0.01*	7.20 ± 0.01*	7.12 ± 0.02*	7.17 ± 0.03*
v	7.35 ± 0.00	7.32 ± 0.01	7.22 ± 0.01*	7.13 ± 0.01*	7.09 ± 0.01*	7.06 ± 0.01*	7.00 ± 0.02*	7.14 ± 0.03*
Lactate (mmol l⁻¹)								
a	1.1 ± 0.1	1.9 ± 0.2	5.3 ± 0.4*	10.8 ± 0.6*	13.7 ± 0.8*	16.0 ± 0.9*	19.0 ± 1.1*	13.8 ± 1.2*
v	1.0 ± 0.1	2.2 ± 0.2	5.4 ± 0.5*	10.8 ± 0.6*	13.4 ± 0.7*	15.6 ± 0.9*	18.6 ± 1.1*	13.6 ± 1.2*
Glucose (mmol l⁻¹)								
a	5.4 ± 0.2	5.3 ± 0.2	5.4 ± 0.2	5.3 ± 0.3	5.2 ± 0.3	5.2 ± 0.3	5.3 ± 0.2	7.3 ± 0.3*
v	5.3 ± 0.2	5.3 ± 0.2	5.3 ± 0.2	5.3 ± 0.2	5.2 ± 0.3	5.2 ± 0.2	5.2 ± 0.2	7.2 ± 0.3*

Values are means ± s.e.m. for 8 subjects. a, arterial. v, right atrium. * Different from rest, $P < 0.05$. P_{O_2} , P_{CO_2} and pH values were corrected for changes in blood temperature.

reached a plateau (range 0.75–3 min) and then declined ($P < 0.05$). No difference in dP/dt_{max} was observed at peak SV and exhaustion in either INC (2287 ± 96 versus 2419 ± 77 mmHg s⁻¹; $P = 0.35$) or CON (2119 ± 271 versus 2193 ± 230 mmHg s⁻¹; $P = 0.72$).

In both INC and CON, there was an increase in haemoglobin concentration, resulting in an elevated arterial O₂ content, despite declining arterial O₂ tension and saturation (Tables 1 and 2; Fig. 2). In INC, systemic O₂ delivery increased linearly to 80% of peak power ($r^2 = 0.998$; $P < 0.001$) and then levelled off (Figs 2–4). In CON, systemic O₂ delivery increased to reach a maximal value of 5.7 ± 0.2 l min⁻¹ after 4.4 ± 0.4 min (range 3–5 min) and then declined 0.35 ± 0.08 l min⁻¹ before exhaustion ($P < 0.05$). During both INC and CON, systemic a–v O₂ difference and O₂ extraction increased until exhaustion. At exhaustion in INC and CON, systemic O₂ extraction was $84 \pm 2\%$ and $87 \pm 1\%$, respectively. In INC, \dot{V}_{O_2} increased linearly to exhaustion and therefore \dot{V}_{O_2max} was reached during the last ~0.5 min ($r^2 = 0.998$; $P < 0.001$; Fig. 3). In CON, \dot{V}_{O_2max} was reached within 4–6 min, maintained for 2.0 ± 0.3 min, but declined 0.14 ± 0.05 l min⁻¹ before exhaustion ($P < 0.05$).

Leg haemodynamics, O₂ transport and O₂ uptake

The rate of increase in LBF and O₂ delivery during incremental exercise was attenuated at intensities above 50% of peak power, reaching a plateau at 73–88% (Fig. 5). The levelling off in LBF, associated with a plateau in leg vascular conductance, attenuated the increase in leg \dot{V}_{O_2} (8.8 ± 0.5 versus 11.9 ± 0.7 ml W⁻¹ min⁻¹ at exhaustion compared to 50% peak power, respectively; $P = 0.003$), despite increasing O₂ extraction. At the systemic level, \dot{Q} reached a plateau at ~80% peak power leading to the blunting of the rate of increase in O₂ delivery per litre of \dot{V}_{O_2} (Fig. 6). In contrast to incremental cycling, LBF, O₂ delivery and leg \dot{V}_{O_2} increased linearly from rest to exhaustion during one-legged knee-extensor exercise ($r^2 = 0.994$ – 0.999 ; $P < 0.0002$), thereby allowing the maintenance of constant LBF, leg O₂ delivery and leg \dot{V}_{O_2} per unit of work (slopes = 75.6, 16.3 and 13.4 ml W⁻¹ min⁻¹, respectively). As depicted in Fig. 7, LBF and leg \dot{V}_{O_2} were higher during maximal cycling compared to knee-extensor exercise. However, leg \dot{V}_{O_2} was lower during cycle compared to knee-extensor exercise when expressed per unit of work (8.8 ± 0.5 versus 12.2 ± 0.6 ml W⁻¹ min⁻¹, respectively;

$P < 0.01$) or estimated active muscle mass (175 ± 9 versus 443 ± 34 ml kg⁻¹ min⁻¹, respectively; $P < 0.01$) due largely to the lower blood flow.

Discussion

This study employed two-legged cycling and one-legged knee-extension as exercise models for investigating whether O₂ transport poses a limitation to aerobic power and capacity in humans. The key observations were: (1) during incremental cycling, \dot{Q} , LBF and O₂ delivery reached a plateau at intensities below $\dot{V}_{O_2\max}$, accompanying a decline in SV and increases in central venous pressure and MAP, (2) with a blunted LBF, leg \dot{V}_{O_2} per unit of work declined despite the increasing O₂ extraction, (3) when LBF increased linearly during knee-extensor exercise, \dot{V}_{O_2} per unit of work was unaltered, (4) during constant load cycling, SV, \dot{Q} , and systemic O₂ delivery and \dot{V}_{O_2} dropped before exhaustion, despite increasing or stable central venous pressure, MAP and

O₂ extraction, and (5) the impaired O₂ supply during cycling was due to a fall in SV, as both the arterial O₂ content and HR continued to increase. These results indicate that an inability of the circulatory system to sustain a linear increase in O₂ delivery to the locomotor limb muscles markedly restrains aerobic power. Moreover, the impaired SV and O₂ delivery during cycling together with the higher muscle \dot{V}_{O_2} with unrestricted circulation during knee-extensor exercise support a preponderant central limitation to aerobic power and capacity in humans.

During incremental cycling exercise, \dot{Q} and systemic O₂ delivery increased linearly to ~80% of peak power but levelled off thereafter. Similarly, the initial linear increase in LBF and O₂ delivery was attenuated at intensities above 50% of peak power, reaching a plateau at 73–88%. The tight 1 : 1 relationship between systemic and locomotor limb O₂ delivery versus \dot{V}_{O_2} at low and moderate exercise intensities was therefore blunted at intensities above ~80% of peak power. This novel finding refutes the theory




Figure 5. Leg haemodynamics during incremental exercise to exhaustion

Leg blood flow, mean arterial pressure, leg vascular conductance, leg O₂ delivery, leg a-v O₂ difference and leg \dot{V}_{O_2} during incremental exercise to exhaustion plotted against the relative increase in power. Data are means \pm S.E.M. for 10 subjects.

that \dot{V}_{O_2} delivery is linearly related to \dot{V}_{O_2} from rest to $\dot{V}_{O_2,max}$. This concept is based on linear regression analysis of: (1) \dot{Q} data that were not normalized (Asmussen & Nielsen, 1952; Chapman *et al.* 1960; Bevegård *et al.* 1963; Åstrand *et al.* 1964; Saltin, 1964; Grimsby *et al.* 1966), (2) haemodynamic data from longitudinal studies on human subjects undergoing changes in physical activity levels (Saltin *et al.* 1968), (3) cross-sectional data from subjects with different training status with the focus of the analysis on whether \dot{Q} could explain the differences in $\dot{V}_{O_2,max}$ between well-trained and sedentary people (Ekblohm & Hermansen, 1968; Ekblohm, 1968), and (4) haemodynamic data in untrained humans who might have failed to attain maximal levels of exertion (Poliner *et al.* 1980; Higginbotham *et al.* 1986). Although these studies collectively documented a tight relationship between \dot{Q} and \dot{V}_{O_2} over a wide range of aerobic capacities, they provide neither insight into the relationship between systemic O_2 delivery and \dot{V}_{O_2} close to maximal exercise, nor specific information on the contribution of locomotor limb muscle O_2 transport to $\dot{V}_{O_2,max}$. The present findings in healthy trained subjects show that systemic and locomotor limb blood flow and O_2 delivery are linearly related to \dot{V}_{O_2} up to 50–90% of $\dot{V}_{O_2,max}$, levelling off before $\dot{V}_{O_2,max}$ is reached. The critical consequence of the plateau in locomotor limb O_2 delivery is the attenuation in the rate of rise in \dot{V}_{O_2} despite increasing O_2 extraction.

Restrictions in systemic and locomotor limb muscle O_2 transport pose a more important limitation to $\dot{V}_{O_2,max}$ and maximal endurance capacity than suspected. In the exercising legs, \dot{V}_{O_2} per unit of work declined from $11.9 \pm 0.7 \text{ ml W}^{-1} \text{ min}^{-1}$ at 50% peak power to $8.8 \pm 0.5 \text{ ml W}^{-1} \text{ min}^{-1}$ at exhaustion, suggesting that the concomitant exponential rise in leg lactate release accompanied a suppression in skeletal muscle aerobic ATP production. The estimate that leg O_2 delivery and \dot{V}_{O_2} would have been $\sim 52\%$ higher (i.e. both $\sim 11 \text{ min}^{-1}$) during cycling exercise if LBF increased linearly until exhaustion, illustrates the magnitude of the blunting of LBF and its effect on locomotor muscle \dot{V}_{O_2} . Limitations in diffusive O_2 transport from the muscle capillary to the mitochondrial cytochrome and/or oxidative capacity of mitochondria could also be restricting muscle \dot{V}_{O_2} based on the observation that leg O_2 extraction is not maximal at exhaustion (Roca *et al.* 1989). Yet, an increase in leg O_2 extraction from the measured value of 87% to a hypothetical 100% would only increase leg \dot{V}_{O_2} by 16% (i.e. $\sim 0.31 \text{ min}^{-1}$), although this is an overestimation given that $\sim 20\%$ of the leg consists of non-muscle tissues with lower O_2 extraction than contracting muscle.

A critical question is whether the leg muscles can indeed increase \dot{V}_{O_2} above the levels observed during maximal cycling exercise. An approach to answer this question is to determine whether quadriceps \dot{V}_{O_2} is elevated when systemic O_2 transport is not limiting during maximal

one-legged knee-extensor exercise (Andersen & Saltin, 1985). In contrast to cycling and in agreement with published reports (Andersen & Saltin, 1985; Richardson *et al.* 1993), leg \dot{V}_{O_2} increased linearly during knee-extensor exercise to exhaustion in parallel with the rise in LBF. Leg \dot{V}_{O_2} per unit of work at fatigue was therefore higher during knee-extensor than cycling exercise (Fig. 7). Moreover, assuming that the quadriceps femoris and all leg muscles are active during knee-extensor and cycling exercise, we estimated that \dot{V}_{O_2} per kilogram of muscle was 3-fold higher during knee-extensor than cycling exercise (Richardson & Saltin 1998). Despite




Figure 6. Relationship between blood flow, vascular conductance and O_2 delivery and \dot{V}_{O_2}

Cardiac output and 2-legged blood flow, systemic and 2-legged vascular conductance, and systemic and 2-legged O_2 delivery plotted against the increases in \dot{V}_{O_2} during incremental exercise to exhaustion. Data are means \pm s.e.m. for 6 subjects.

the uncertainty of the assumption, it is clear that the quadriceps muscles as the only muscles generating power during knee-extensor exercise, are consuming more O_2 (i.e. 1.3 l min^{-1} ; 103 W ; $\sim 2.9 \text{ kg}$) than they are as mere contributors to power generation during cycling (i.e. 2.0 l min^{-1} ; 225 W ; $\sim 11.4 \text{ kg}$). Collectively, these observations reveal that during maximal whole body exercise: (1) locomotor muscle \dot{V}_{O_2} and exercise endurance could be improved if blood flow increased linearly, (2) the rates of mitochondrial oxidation and O_2 transport from capillary to mitochondrial cytochrome are not maximal, and (3) convective O_2 transport to contracting skeletal muscle fibres is severely restricted owing to the lower blood flow.

During incremental cycling, \dot{Q} and LBF plateaued at $\sim 80\%$ peak power in parallel to a marked blunting of systemic and leg vascular conductance, indicating an enhanced sympathetic vasoconstrictor activity. This agrees with the plateau in LBF at high cycling intensities in humans (Knight *et al.* 1992; Rosenmeier *et al.* 2004) and the plateau in \dot{Q} and skeletal muscle blood flow before exhaustion in miniature swine running on a treadmill (Armstrong *et al.* 1987), but contrasts with the linear increase in LBF during one-legged knee-extensor exercise in humans when the systemic circulation is not compromised (Andersen & Saltin, 1985; Richardson *et al.* 1993). It therefore seems that during exercise with a large muscle mass, reflexes signalling the plateau in \dot{Q} override the local vasodilatory stimuli responsible for the partial or full linear increase in LBF with incremental cycling and knee-extensor exercise, respectively. In support of this, a human study showed that the blunting of LBF and vascular

conductance at high cycling intensities is associated with an exponential rise in circulating noradrenaline outstripping the increase of the vasodilator ATP (Rosenmeier *et al.* 2004). It is likely that the upper body limb and postural muscles as well as the heart and respiratory muscles become more active, thereby contributing to the linear increase in systemic \dot{V}_{O_2} above 80% peak power. This scenario implies that the upper body muscles and organs are competing with the exercising legs for the available \dot{Q} (Harms *et al.* 1997, 1998). However, the redistribution of blood flow to upper body muscles and organs ought to be small as the plateau in \dot{Q} accounts for the majority of the LBF response. On the other hand, perfusion pressure did not reduce LBF as MAP increased until exhaustion. Consequently, the suppression in blood flow to the exercising legs during cycling appears to be, for the most part, the result of the blunted \dot{Q} and the overriding sympathetic vasoconstrictor activity to the muscle microvasculature (Pawelczyk *et al.* 1992).

The similar maximal haemodynamic responses and blunting of O_2 transport before exhaustion during INC and CON provide further insight into the limits of cardiovascular regulation in exercising humans. Even though $\dot{V}_{O_{2\max}}$ was maintained over a longer period during constant load cycling, both types of maximal exercise were characterized by essentially the same peak \dot{Q} (27 and 28 l min^{-1} for INC and CON, respectively), HR (190 and $192 \text{ beats min}^{-1}$), MAP (129 and 136 mmHg), systemic O_2 delivery (5.7 and 5.8 l min^{-1}), systemic O_2 extraction (84 and 87%) and $\dot{V}_{O_{2\max}}$ (4.8 l min^{-1}). This indicates that the limits of the cardiovascular system and a true $\dot{V}_{O_{2\max}}$ were reached during both types of cycling. Five




Figure 7. Quadriceps muscle perfusion and \dot{V}_{O_2} during maximal knee-extensor and cycling exercise
Maximal values for leg blood flow and \dot{V}_{O_2} during maximal knee-extensor and cycling exercise expressed as absolute values (A), in relation to peak power (B; 103 ± 4 and $225 \pm 10 \text{ W}$, respectively), and in relation to the estimated active muscle mass (C), assuming that all quadriceps muscles (2.9 kg) and all leg muscles (11.4 kg) were recruited during maximal knee-extensor and cycling exercise, respectively. Data are means \pm s.e.m. for 10 subjects. * Different from knee-extensor exercise, $P < 0.05$.

decades ago, Mitchell *et al.* (1958) provided data on the determinants of $\dot{V}_{O_{2max}}$. Arguing in favour of a limiting \dot{Q} , they found a lower \dot{Q} during supramaximal compared to maximal exercise (18.2 *versus* 21.0 l min⁻¹, respectively), but $\dot{V}_{O_{2max}}$ was the same (2.81 *versus* 2.87 l min⁻¹; $n = 6$) because of the widening of the systemic a–v O₂ difference. Our study extends previous work by simultaneously looking at the dynamics of the central and exercising limb circulations, allowing an assessment of the contribution of the locomotor muscle to $\dot{V}_{O_{2max}}$. Because LBF and \dot{V}_{O_2} are impaired during maximal constant load cycling (González-Alonso & Calbet, 2003), we surmise that the locomotor skeletal muscles are the main tissue accounting for the restrictions in peripheral blood flow and O₂ delivery during INC and CON.

Within the central circulation, the impaired O₂ supply was associated with a fall in SV (20–27 ml beat⁻¹), as both arterial O₂ content and HR continued to increase. This is congruent with reports during constant and incremental exercise showing a decline in SV (Keul *et al.* 1981; Higginbotham *et al.* 1986; Spina *et al.* 1992; Seals *et al.* 1994; Proctor *et al.* 1998; McCole *et al.* 1999; González-Alonso & Calbet, 2003; González-Alonso *et al.* 2004), but contrasts with the bulk of studies using incremental exercise showing either a plateau or an increase in SV with continuously increasing \dot{Q} from moderate to peak exercise (Åstrand *et al.* 1964; Poliner *et al.* 1980; Rubal *et al.* 1986; Spina *et al.* 1992; Gledhill *et al.* 1994; Seals *et al.* 1994; Fleg *et al.* 1994; Proctor *et al.* 1998). Differences in exercise protocols, levels of exertion, exercise mode, sex, age and training status might account for the discrepancy in the SV responses.

The decline in SV described here could be attributed to alterations in cardiac preload, left ventricular afterload and/or left ventricular contractility (Rowell, 1974, 1993; Poliner *et al.* 1980; Higginbotham *et al.* 1986). However, a decline in preload does not seem to be a factor because in both trials central venous pressure continued to increase until exhaustion. Enhanced afterload might not be an important factor either, since SV increased early in exercise in parallel with increases in systolic blood pressure and MAP in both INC and CON and declined to a greater extent during CON when systolic blood pressure and MAP were maintained. Lastly, a depression in left ventricular contractility reducing SV is at odds with the finding that dP/dt_{max} at peak SV and at exhaustion were not different. The rate–pressure product of HR and MAP increased until exhaustion in both maximal tests, indicating that myocardial O₂ demand was rising when SV declined. In this setting, an increase in myocardial \dot{V}_{O_2} can only occur by an increase in O₂ delivery provided by augmented coronary blood flow because the O₂ extraction reserve is minimal. The impaired circulation and aerobic energy turnover in skeletal muscle raises the daunting possibility that alterations in cardiac metabolism contribute to the SV decline.

Alternatively, the observation that SV declined at a HR of 170–180 beats min⁻¹ during both INC and CON raises the possibility that severe tachycardia reduces SV. Studies in humans and dogs manipulating HR by pacing the heart demonstrate that severe tachycardia leads to disproportional reductions in diastolic filling time and left ventricular end-diastolic volume which compromise SV and \dot{Q} (Templeton *et al.* 1972; Weisfeldt *et al.* 1978; Parke & Case, 1979; Sheriff *et al.* 1993). Consistent with the increase in core temperature to 39–40°C, human studies demonstrate that hyperthermia-induced tachycardia reduces SV during exercise (Fritzsche *et al.* 1999; González-Alonso *et al.* 1997, 1999, 2000a) and that blunting core hyperthermia and HR restores most of the fall in $\dot{V}_{O_{2max}}$ evoked by heat stress (Nybo *et al.* 2001). Although studies independently altering HR and directly examining cardiac circulation, metabolism and function are required to determine the mechanism, it seems that the decline in SV during maximal exercise is related, at least in part, to the restriction in left ventricular filling time and left ventricular end-diastolic volume that accompanies severe tachycardia and hyperthermia.

In summary, the present findings in trained humans show that systemic and locomotor limb O₂ delivery does not increase linearly from rest to $\dot{V}_{O_{2max}}$, but plateaus at intensities below $\dot{V}_{O_{2max}}$, resulting in the blunting of locomotor limb \dot{V}_{O_2} despite the increasing O₂ extraction. Similarly, systemic O₂ delivery and \dot{V}_{O_2} decline during constant load cycling despite the increasing O₂ extraction. In both types of maximal exercise, the impaired systemic O₂ delivery was associated with a decline in SV. The attenuation in LBF blunting leg O₂ delivery and \dot{V}_{O_2} during incremental cycling appears to be largely related to the plateau in \dot{Q} and an enhanced muscle sympathetic vasoconstrictor activity. In contrast to two-legged cycling, LBF and \dot{V}_{O_2} increased until volitional exhaustion during one-legged knee-extensor exercise when O₂ transport was not limited. Collectively, these findings support the hypothesis that restrictions in O₂ supply to locomotor limb muscles impose a limitation to aerobic power and capacity in humans.

References

- Andersen P & Saltin B (1985). Maximal perfusion of skeletal muscle in man. *J Physiol* **366**, 233–249.
- Armstrong RB, Delp MD, Goljan EF & Laughlin MH (1987). Distribution of blood flow in muscles of miniature swine during exercise. *J Appl Physiol* **62**, 1285–1298.
- Asmussen E & Nielsen M (1952). The cardiac output in rest and work determined simultaneously by the acetylene and the dye injection methods. *Acta Physiol Scand* **27**, 217–230.
- Åstrand PO (1952). *Experimental Studies of Physical Working Capacity in Relation to Sex and Age*. Munksgaard, Copenhagen.

- Åstrand PO, Cuddy TE, Saltin B & Stenberg J (1964). Cardiac output during submaximal and maximal work. *J Appl Physiol* **19**, 268–274.
- Åstrand PO & Saltin B (1961). Oxygen uptake during the first minutes of heavy muscular exercise. *J Appl Physiol* **16**, 971–976.
- Barratt-Boyes BG & Wood EH (1956). The oxygen saturation of blood in the venae cavae, right-heart chambers, and pulmonary vessels of healthy subjects. *J Lab Clin Med* **50**, 93–106.
- Bevegård S, Holmgren A & Johnsson B (1963). Circulatory studies in well-trained athletes at rest and during heavy exercise, with special reference to stroke volume and the influence of body composition. *Acta Physiol Scand* **57**, 26–50.
- Chapman CB, Fisher JN & Sproule BJ (1960). Behavior of stroke volume at rest and during exercise in human beings. *J Clin Invest* **39**, 1208–1213.
- Day JR, Rossiter HB, Coats EM, Skasick A & Whipp BJ (2003). The maximally attainable $\dot{V}O_2$ during exercise in humans: the peak vs. maximum issue. *J Appl Physiol* **95**, 1901–1907.
- Eklblom B (1968). Effect of physical training on oxygen transport system in man. *Acta Physiol Scand Suppl* **328**, 1–45.
- Eklblom B & Hermansen L (1968). Cardiac output in athletes. *J Appl Physiol* **25**, 619–625.
- Ekelund LG & Holmgren A (1967). Central hemodynamics during exercise. *Circ Res* **21** (Suppl. 1), I33–I43.
- Fleg JL, Schulman SP, O'Connor FC, Gerstenblith G, Becker LC, Fortney S, Goldberg AP & Lakatta EG (1994). Cardiovascular responses to exhaustive upright cycle exercise in highly trained older men. *J Appl Physiol* **77**, 1500–1506.
- Fritzsche RG, Switzer TW, Hodgkinson BJ & Coyle EF (1999). Stroke volume decline during prolonged exercise is influenced by the increase in heart rate. *J Appl Physiol* **86**, 799–805.
- Froelicher VF, Brammell H, Davis G, Noguera I, Stewart A & Lancaster MC (1972). A comparison of three maximal treadmill exercise protocols. *J Appl Physiol* **36**, 720–725.
- Gledhill N, Cox D & Jamnik R (1994). Endurance athletes' stroke volume does not plateau: major advantage is diastolic function. *Med Sci Sports Exerc* **26**, 1116–1121.
- González-Alonso J & Calbet J (2003). Reductions in systemic and skeletal muscle blood flow and oxygen delivery limit maximal aerobic capacity in humans. *Circulation* **107**, 824–830.
- González-Alonso J, Dalsgaard MK, Osada T, Volianitis S, Dawson EA, Yoshiga CC & Secher NH (2004). Brain and central haemodynamics and oxygenation during maximal exercise in humans. *J Physiol* **557**, 331–342.
- González-Alonso J, Mora-Rodríguez R & Coyle EF (1997). Dehydration markedly impairs cardiovascular function in hyperthermic endurance athletes during exercise. *J Appl Physiol* **82**, 1229–1236.
- González-Alonso J, Mora-Rodríguez R & Coyle EF (2000a). Stroke volume during exercise: interaction of environment and hydration. *Am J Physiol* **278**, H321–H330.
- González-Alonso J, Quistorff B, Krstrup P, Bangsbo J & Saltin B (2000b). Heat production in human skeletal muscle at onset of intense dynamic exercise. *J Physiol* **524**, 603–615.
- González-Alonso J, Teller C, Andersen SL, Jensen FB, Hyldig T & Nielsen B (1999). Influence of body temperature on the development of fatigue during prolonged exercise in the heat. *J Appl Physiol* **86**, 1032–1039.
- Grimby G, Nilsson NJ & Saltin B (1966). Cardiac output during submaximal and maximal exercise in active middle-aged athletes. *J Appl Physiol* **21**, 1150–1156.
- Harms CA, Babcock MA, McClaran SR, Pegelow DF, Nickle GA, Nelson WB & Dempsey JA (1997). Respiratory muscle work compromises leg blood flow during maximal exercise. *J Appl Physiol* **82**, 1573–1583.
- Harms CA, Wetter TJ, McClaran SR, Pegelow DF, Nickle GA, Nelson WB, Hanson P & Dempsey JA (1998). Effects of respiratory muscle work on cardiac output and its distribution during maximal exercise. *J Appl Physiol* **85**, 609–618.
- Higginbotham MB, Morris KG, Williams RS, McHale PA, Coleman RE & Cobb FR (1986). Regulation of stroke volume during submaximal and maximal upright exercise in normal man. *Circ Res* **58**, 281–291.
- Hill AV & Lupton H (1923). Muscular exercise, lactic acid, and the supply and utilization of oxygen. *Q J Med* **16**, 135–171.
- Holmgren A (1956). Circulatory changes during muscular work in man: with special reference to arterial and central venous pressures in the systemic circulation. *Scand J Clin Lab Invest* **8** (Suppl. 24), 1–97.
- Keul J, Dickhuth HH, Simon G & Lehmann M (1981). Effect of static and dynamic exercise on heart volume, contractility, and left ventricular dimensions. *Circ Res* **48**, I162–I170.
- Knight DR, Poole DC, Schaffartzik W, Gay HJ, Prediletto R, Hogan MC & Wagner P (1992). Relationship between body and leg $\dot{V}O_2$ during maximal cycle ergometry. *J Appl Physiol* **73**, 1114–1121.
- McCole SD, Brown MD, Moore GE, Zmuda JM, Cwynar JD & Hagberg JM (1999). Cardiovascular hemodynamics with increasing exercise intensities in postmenopausal women. *J Appl Physiol* **87**, 2334–2340.
- Meyers J, Walsh D, Sullivan M & Froelicher V (1990). Effect of sampling on variability and plateau in oxygen uptake. *J Appl Physiol* **68**, 404–410.
- Mitchell JH, Sproule BJ & Chapman CB (1958). The physiological meaning of the maximal oxygen intake test. *J Clin Invest* **37**, 538–547.
- Nybo L, Jensen T, Nielsen B & González-Alonso J (2001). Effects of hyperthermia with and without dehydration on $\dot{V}O_2$ kinetics during intense exercise. *J Appl Physiol* **90**, 1057–1064.
- Parke JO & Case RB (1979). Normal left ventricular function. *Circulation* **60**, 4–12.
- Pawelczyk JA, Hanel B, Pawelczyk RA, Warberg J & Secher NH (1992). Leg vasoconstriction during dynamic exercise with reduced cardiac output. *J Appl Physiol* **73**, 1838–1846.
- Poliner LR, Dehmer GJ, Lewis SE, Parkey RW, Blomqvist CG & Willerson JT (1980). Left ventricular performance in normal subjects: a comparison of the responses to exercise in the upright and supine positions. *Circulation* **62**, 528–534.
- Pollock ML, Bohannon RL, Cooper KH, Ayres JJ, Ward A, White SR & Linnerud AC (1976). A comparative analysis of four protocols for maximal treadmill stress testing. *Am Heart J* **92**, 39–46.

- Proctor DN, Beck KC, Shen PH, Eickhoff TJ, Halliwill JR & Joyner MJ (1998). Influence of age and gender on cardiac output-VO₂ relationships during submaximal cycle ergometry. *J Appl Physiol* **84**, 599–605.
- Reeves JT, Groves BM, Cymerman A, Sutton JR, Wagner PD, Turkevich D & Houston CS (1990). Operation Everest II: cardiac filling pressures during cycle exercise at sea level. *Resp Physiol* **80**, 147–154.
- Richardson RS, Poole DC, Knight DR, Kurdak SS, Hogan MC, Grassi B, Johnson EC, Kendrick KF, Erickson BK & Wagner PD (1993). High muscle blood flow in man: is maximal O₂ extraction compromised? *J Appl Physiol* **75**, 1911–1926.
- Richardson RS & Saltin B (1998). Human muscle blood flow and metabolism studied in the isolated quadriceps muscles. *Med Sci Sports Exerc* **30**, 28–33.
- Roca J, Hogan MC, Story D, Bebout DE, Haab P, Gonzalez R, Ueno O & Wagner PD (1989). Evidence for tissue diffusion limitation of VO₂max in normal humans. *J Appl Physiol* **67**, 291–299.
- Rosenmeier JB, Hansen J & González-Alonso J (2004). Circulating ATP-induced vasodilatation overrides sympathetic vasoconstrictor activity in human skeletal muscle. *J Physiol* **558**, 351–365.
- Rowell LB (1974). Human cardiovascular adjustments to exercise and thermal stress. *Physiol Rev* **54**, 75–159.
- Rowell LB (1993). *Human Cardiovascular Control*. Oxford University Press, New York.
- Rowell LB, O'Leary DS & Kellogg DL Jr (1996). Integration of cardiovascular control systems in dynamic exercise. In *Handbook of Physiology*, section 12, *Exercise: Regulation and Integration of Multiple Systems*, chap. 17, pp. 770–838. American Physiological Society, Bethesda, MD, USA.
- Rowell LB, Taylor HL & Wang Y (1964). Limitations to prediction of maximal oxygen intake. *J Appl Physiol* **19**, 919–927.
- Rubal BJ, Moody JM, Damore S, Bunker SR & Diaz NM (1986). Left ventricular performance of the athletic heart during upright exercise: a heart rate-controlled study. *Med Sci Sports Exerc* **18**, 134–140.
- Saltin B (1964). Circulatory response to submaximal and maximal exercise after thermal dehydration. *J Appl Physiol* **19**, 1125–1132.
- Saltin B, Blomqvist G, Mitchell JH, Johnson RL Jr, Wildenthal K & Chapman CB (1968). Response to exercise after bed rest and after training. *Circulation* **38** (5 Suppl.), VIII–78.
- Seals DR, Hagberg JM, Spina RJ, Rogers MA, Schechtman KB & Ehsani AA (1994). Enhanced left ventricular performance in endurance trained older men. *Circulation* **89**, 198–205.
- Sheriff DD, Zhou XP, Scher AM & Rowell LB (1993). Dependence of cardiac filling pressure on cardiac output during rest and dynamic exercise in dogs. *Am J Physiol* **265**, H316–H322.
- Spina RJ, Ogawa T, Martin WH 3rd, Coggan AR, Holloszy JO & Ehsani AA (1992). Exercise training prevents decline in stroke volume during exercise in young healthy subjects. *J Appl Physiol* **72**, 2458–2462.
- Taylor HL, Buskirk E & Henschel A (1955). Maximal oxygen intake as an objective measure of cardio-respiratory performance. *J Appl Physiol* **8**, 73–80.
- Templeton GH, Ecker RR & Mitchel JH (1972). Left ventricular stiffness during diastole and systole: the influence of changes in volume and inotropic state. *Cardiovasc Res* **6**, 95–100.
- Weisfeldt ML, Frederiksen JW, Yin FCP & Weiss JL (1978). Evidence of incomplete left ventricular relaxation in the dog. *J Clin Invest* **62**, 1296–1302.
- Whipp BJ & Wasserman K (1972). Oxygen kinetics for various intensities of constant-load exercise. *J Appl Physiol* **33**, 351–356.

Acknowledgements

We give special thanks to the volunteer subjects for their enthusiasm. We also thank Peter Nissen, Troels Munch and Jacob Mørkeberg for the excellent technical assistance. This study was supported by the Gatorade Sports Science Institute and the Novo Nordisk Foundation. J.G.-A. and N.H.S. were supported by The Copenhagen Hospital System.

Limitations to systemic and locomotor limb muscle oxygen delivery and uptake during maximal exercise in humans

Stefan P Mortensen, Ellen A Dawson, Chie C Yoshiga, Mads K Dalsgaard, Rasmus Damsgaard, Niels H Secher and José González-Alonso

J. Physiol. 2005;566;273-285; originally published online Apr 28, 2005;

DOI: 10.1113/jphysiol.2005.086025

This information is current as of October 11, 2007

**Updated Information
& Services**

including high-resolution figures, can be found at:
<http://jp.physoc.org/cgi/content/full/566/1/273>

Permissions & Licensing

Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at:
<http://jp.physoc.org/misc/Permissions.shtml>

Reprints

Information about ordering reprints can be found online:
<http://jp.physoc.org/misc/reprints.shtml>