

**You might find this additional information useful...**

---

This article cites 61 articles, 42 of which you can access free at:

<http://jap.physiology.org/cgi/content/full/100/2/744#BIBL>

This article has been cited by 9 other HighWire hosted articles, the first 5 are:

**Cardiac output and leg and arm blood flow during incremental exercise to exhaustion on the cycle ergometer**

J. A. L. Calbet, J. Gonzalez-Alonso, J. W. Helge, H. Sondergaard, T. Munch-Andersen, R. Boushel and B. Saltin

*J Appl Physiol*, September 1, 2007; 103 (3): 969-978.

[Abstract] [Full Text] [PDF]

**Oxygen delivery by blood determines the maximal VO<sub>2</sub> and work rate during whole body exercise in humans: in silico studies**

P. Liguzinski and B. Korzeniewski

*Am J Physiol Heart Circ Physiol*, July 1, 2007; 293 (1): H343-H353.

[Abstract] [Full Text] [PDF]

**Continued divergence in VO<sub>2</sub> max of rats artificially selected for running endurance is mediated by greater convective blood O<sub>2</sub> delivery**

N. C. Gonzalez, S. D. Kirkton, R. A. Howlett, S. L. Britton, L. G. Koch, H. E. Wagner and P. D. Wagner

*J Appl Physiol*, November 1, 2006; 101 (5): 1288-1296.

[Abstract] [Full Text] [PDF]

**Is the healthy respiratory system (always) built for exercise?**

J. A. Dempsey

*J. Physiol.*, October 15, 2006; 576 (2): 339-340.

[Full Text] [PDF]

**Effects of ATP-induced leg vasodilation on VO<sub>2</sub> peak and leg O<sub>2</sub> extraction during maximal exercise in humans**

J. A. L. Calbet, C. Lundby, M. Sander, P. Robach, B. Saltin and R. Boushel

*Am J Physiol Regulatory Integrative Comp Physiol*, August 1, 2006; 291 (2): R447-R453.

[Abstract] [Full Text] [PDF]

Medline items on this article's topics can be found at <http://highwire.stanford.edu/lists/artbytopic.dtl> on the following topics:

Physiology .. Hemodynamics  
Physiology .. Blood Circulation  
Medicine .. Cardiac Output  
Medicine .. Fitness (Physical Activity)  
Medicine .. Exercise  
Medicine .. Exertion

Updated information and services including high-resolution figures, can be found at:

<http://jap.physiology.org/cgi/content/full/100/2/744>

Additional material and information about *Journal of Applied Physiology* can be found at:

<http://www.the-aps.org/publications/jappl>

---

This information is current as of October 12, 2007 .

## Point: In health and in a normoxic environment, $\dot{V}O_{2\max}$ is limited primarily by cardiac output and locomotor muscle blood flow

Starting in the 1950s, a number of experiments provided the experimental evidence supporting the original concept elaborated on by Hill and Lupton (12): in health,  $\dot{V}O_{2\max}$  in normoxia is limited primarily by cardiac output and locomotor muscle blood flow (17). The main variable accounting for the difference in  $\dot{V}O_{2\max}$  between sedentary subjects and athletes is maximal cardiac output, such that a linear relationship was observed between  $\dot{V}O_{2\max}$  and maximal cardiac output, showing that 5.9–7.5 l/min of cardiac output is needed per liter of  $\dot{V}O_{2\max}$  (5, 10, 17, 26). Part of the variability in the relationship between  $\dot{V}O_{2\max}$  and cardiac output was attributed to the variation in hemoglobin concentration, with a smaller contribution of the systemic a-v difference (5, 10, 17, 26). It was also shown that maximal exercise stroke volume was the main factor explaining the differences between subjects in maximal cardiac output (5, 10, 17, 26). A cause and effect relationship between oxygen delivery and  $\dot{V}O_{2\max}$  has been established by showing that experimental interventions increasing oxygen delivery are accompanied by an elevation of  $\dot{V}O_{2\max}$  and vice versa (6, 16).

All experimental procedures causing a reduction of maximal cardiac output are associated with a lower  $\dot{V}O_{2\max}$ . Reducing blood volume is associated with lower maximal cardiac output and  $\dot{V}O_{2\max}$  (16). Bed rest studies showed that the main factor accounting for the reduction in  $\dot{V}O_{2\max}$  was the lower maximal cardiac output attained after bed rest (27), because maximal exercise  $O_2$  fractional extraction is close to 90% after bed rest. Treatment with beta-blockers is accompanied by a reduction of maximal cardiac output and leg blood flow, which accounts for most of the reduction observed in  $\dot{V}O_{2\max}$  (21). The  $CaO_2$  may be reduced by reducing hemoglobin concentration isovolemically and by carbon monoxide administration. These two interventions show a reduction in  $\dot{V}O_{2\max}$  that is proportional to the magnitude of the reduction achieved in  $CaO_2$  (6, 15, 23, 30).

The influence of locomotor muscle oxygen delivery for  $\dot{V}O_{2\max}$  in trained and untrained muscles was studied in the 1970s (3, 8, 28). With the use of a one-leg training model (in the cycle ergometer), Gleser (8) reported a 16% improvement of one-leg peak  $\dot{V}O_2$  that was accompanied by a 13% enhancement of the peak cardiac output during incremental exercise with the trained leg. However, neither  $\dot{V}O_{2\max}$  nor maximal cardiac output was enhanced after one-leg training when the exercise test was performed with the two legs. Thus the study by Gleser suggests that the increase in  $\dot{V}O_{2\max}$  was brought about via an enhancement of cardiac output and, likely, leg blood flow. Clausen et al. (3) reported a 10% greater peak  $\dot{V}O_2$  during arm cranking after a period of endurance training with the legs in the cycle ergometer. The increase in arm  $\dot{V}O_2$  was accompanied by 10 and 12% greater mean arterial pressure and peak cardiac output, also suggesting that  $\dot{V}O_{2\text{peak}}$  during exercise with a small muscle mass is limited by locomotor muscle blood flow. In the study by Saltin et al. (28), the subjects that performed one-leg endurance training in the cycle ergometer improved their  $\dot{V}O_{2\max}$  by 24% during an incremental exercise to exhaustion with the trained leg. Interestingly, the contralat-

eral leg that was not submitted to training also improved its  $\dot{V}O_{2\max}$  (6%). However, when the subjects carried out a two-legged incremental exercise the  $\dot{V}O_{2\max}$  was improved only by 11%. Thus the improvement observed during two-leg exercise was a bit less than expected if the limitation to  $\dot{V}O_{2\max}$  had been only of peripheral origin, suggesting that in that study part of the limitation to  $\dot{V}O_{2\max}$  during two-leg exercise is due to insufficient perfusion. A subsequent one-leg training study by Klausen et al. (13) adds further evidence. Their subjects trained each leg on the cycle ergometer individually. After the training, peak leg  $\dot{V}O_2$  during exercise on the cycle ergometer was 16% higher during one-leg than during two-leg exercise, due to a 23% higher peak leg blood flow during one-leg maximal exercise compared with two-leg maximal exercise. In contrast, before training, peak leg  $\dot{V}O_2$  was the same during one-leg cycling compared with two-leg cycling, despite the fact that leg blood flow was 8% higher during one-leg exercise. This study suggests that in the trained state, the dependency of  $\dot{V}O_{2\max}$  on oxygen delivery may be accentuated.

Further evidence for a cause and effect relationship between  $\dot{V}O_{2\max}$  and locomotory muscle oxygen delivery was obtained by Harms et al. (11). They showed that if the respiratory muscles are loaded, exercise capacity and locomotory muscle blood flow and  $\dot{V}O_2$  is reduced, suggesting that maneuvers redistributing part of the blood flow away from the locomotory muscles reduces exercise capacity and  $\dot{V}O_{2\max}$  (11) and vice versa. A similar conclusion was reached by Gonzalez-Alonso and Calbet (9). In their study, subjects performed constant intensity exercise to exhaustion under normothermic and hyperthermic conditions. In both conditions, fatigue was preceded by a reduction of cardiac output and leg blood flow. Moreover, we recently showed that during whole body upright exercise the combined maximal muscular vascular conductances of the limbs outweighs the pumping capacity of the heart in humans, meaning that  $\dot{V}O_{2\max}$  is limited by  $O_2$  delivery. With the use of data from the latter, we estimated that if the human with well-trained leg and arms muscles was able to use the full potential for  $\dot{V}O_2$  of the four limbs, then their  $\dot{V}O_{2\max}$  could be about 20% higher than actually measured (2).

Although  $\dot{V}O_{2\max}$  is a function of locomotor muscle blood flow, this does not exclude the possibility that other mechanisms marginally contribute to achieve  $\dot{V}O_{2\max}$  in normoxia, as, for example, exercise-induced arterial hypoxemia (4, 19), a diffusional limitation between the capillaries and the mitochondria of the active muscle fibers (24), and lower  $O_2$  extraction capacity in some muscles (1). However, in all these conditions, peak  $\dot{V}O_2$  is increased if the limitation is somehow overcome and more  $O_2$  is made available to the mitochondria (6, 14, 22, 25). Thus the bulk of the experimental evidence accumulated during the last 80 years argues in favor of cardiac output and oxygen delivery setting the limit for maximal oxygen uptake in normoxia. All these observations also argue against theories attributing the limitation of  $\dot{V}O_{2\max}$  to brain processes as the “Central Governor Model” during exercise in normoxia carried out by healthy subjects (20). This model postulates that processes arising in the brain itself, triggered or modulated by

sensory feedback, inhibit somehow the central command, causing the exercise to terminate (20). This model has revitalized some ideas brought about more than a century ago, as reviewed by Gandevia (7). However, experimental evidence obtained during exercise with hyperthermia (18) and during exercise in chronic hypoxia (29) demonstrated that, at least during brief efforts aimed at producing a maximal leg or hand grip voluntary contraction, the ability to recruit the motor units is preserved even when measured close to exhaustion.

In summary, in healthy humans,  $\dot{V}O_{2\max}$  at sea level is limited by systemic oxygen delivery and especially by  $O_2$  delivery to the locomotor muscles. Oxygen delivery, in turn, depends on the ability of the cardiorespiratory system (i.e., lungs, heart, and blood) to transport and distribute appropriately  $O_2$  to the active motor units, rather than on the mitochondrial oxidative capacity, which in human skeletal muscles exceeds widely maximal  $O_2$  supply in all known exercise models.

#### REFERENCES

- Calbet JA, Holmberg HC, Rosdahl H, van Hall G, Jensen-Urstad M, and Saltin B. Why do arms extract less oxygen than legs during exercise? *Am J Physiol Regul Integr Comp Physiol* 289: R1448–R1458, 2005.
- Calbet JA, Jensen-Urstad M, Van Hall G, Holmberg HC, Rosdahl H, and Saltin B. Maximal muscular vascular conductances during whole body upright exercise in humans. *J Physiol* 558: 319–331, 2004.
- Clausen JP, Klausen K, Rasmussen B, and Trap-Jensen J. Central and peripheral circulatory changes after training of the arms or legs. *Am J Physiol* 225: 675–682, 1973.
- Dempsey JA and Wagner PD. Exercise-induced arterial hypoxemia. *J Appl Physiol* 87: 1997–2006, 1999.
- Eklblom B and Hermansen L. Cardiac output in athletes. *J Appl Physiol* 25: 619–625, 1968.
- Eklblom B, Wilson G, and Astrand PO. Central circulation during exercise after venesection and reinfusion of red blood cells. *J Appl Physiol* 40: 379–383, 1976.
- Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev* 81: 1725–1789, 2001.
- Gleser MA. Effects of hypoxia and physical training on hemodynamic adjustments to one-legged exercise. *J Appl Physiol* 34: 655–659, 1973.
- Gonzalez-Alonso J and Calbet JA. Reductions in systemic and skeletal muscle blood flow and oxygen delivery limit maximal aerobic capacity in humans. *Circulation* 107: 824–830, 2003.
- Grimby G, Nilsson NJ, and Saltin B. Cardiac output during submaximal and maximal exercise in active middle-aged athletes. *J Appl Physiol* 21: 1150–1156, 1966.
- Harms CA, Babcock MA, McClaran SR, Pegelow DF, Nickle GA, Nelson WB, and Dempsey JA. Respiratory muscle work compromises leg blood flow during maximal exercise. *J Appl Physiol* 82: 1573–1583, 1997.
- Hill AV and Lupton H. Muscular exercise, lactic acid, and the supply and utilization of oxygen. *Q J Med* 16: 135–171, 1923.
- Klausen K, Secher NH, Clausen JP, Hartling O, and Trap-Jensen J. Central and regional circulatory adaptations to one-leg training. *J Appl Physiol* 52: 976–983, 1982.
- Knight DR, Schaffartzik W, Poole DC, Hogan MC, Bebout DE, and Wagner PD. Effects of hyperoxia on maximal leg  $O_2$  supply and utilization in men. *J Appl Physiol* 75: 2586–2594, 1993.
- Koskolou MD, Roach RC, Calbet JA, Radegran G, and Saltin B. Cardiovascular responses to dynamic exercise with acute anemia in humans. *Am J Physiol Heart Circ Physiol* 273: H1787–H1793, 1997.
- Krip B, Gledhill N, Jannik V, and Warburton D. Effect of alterations in blood volume on cardiac function during maximal exercise. *Med Sci Sports Exerc* 29: 1469–1476, 1997.
- Mitchell JH, Sproule BJ, and Chapman CB. The physiological meaning of the maximal oxygen intake test. *J Clin Invest* 37: 538–547, 1958.
- Nielsen B, Hales JR, Strange S, Christensen NJ, Warberg J, and Saltin B. Human circulatory and thermoregulatory adaptations with heat acclimation and exercise in a hot, dry environment. *J Physiol* 460: 467–485, 1993.
- Nielsen HB. Arterial desaturation during exercise in man: implication for  $O_2$  uptake and work capacity. *Scand J Med Sci Sports* 13: 339–358, 2003.
- Noakes TD, St Clair Gibson A, and Lambert EV. From catastrophe to complexity: a novel model of integrative central neural regulation of effort and fatigue during exercise in humans. *Br J Sports Med* 38: 511–514, 2004.
- Pawelczyk JA, Hanel B, Pawelczyk RA, Warberg J, and Secher NH. Leg vasoconstriction during dynamic exercise with reduced cardiac output. *J Appl Physiol* 73: 1838–1846, 1992.
- Richardson RS, Grassi B, Gavin TP, Haseler LJ, Tagore K, Roca J, and Wagner PD. Evidence of  $O_2$  supply-dependent  $\dot{V}O_{2\max}$  in the exercise-trained human quadriceps. *J Appl Physiol* 86: 1048–1053, 1999.
- Roach RC, Koskolou MD, Calbet JA, and Saltin B. Arterial  $O_2$  content and tension in regulation of cardiac output and leg blood flow during exercise in humans. *Am J Physiol Heart Circ Physiol* 276: H438–H445, 1999.
- Roca J, Hogan MC, Story D, Bebout DE, Haab P, Gonzalez R, Ueno O, and Wagner PD. Evidence for tissue diffusion limitation of  $\dot{V}O_{2\max}$  in normal humans. *J Appl Physiol* 67: 291–299, 1989.
- Romer LM, Haverkamp HC, Lovering AT, Pegelow DF, and Dempsey JA. Effect of exercise-induced arterial hypoxemia on quadriceps muscle fatigue in healthy humans. *Am J Physiol Regul Integr Comp Physiol*. First published September 15, 2005 [doi:10.1152/ajpregu.00332.2005].
- Saltin B. Circulatory response to submaximal and maximal exercise after thermal dehydration. *J Appl Physiol* 19: 1125–1132, 1964.
- Saltin B, Blomqvist G, Mitchell JH, Johnson RL Jr, Wildenthal K, and Chapman CB. Response to exercise after bed rest and after training. *Circulation* 38: VIII–78, 1968.
- Saltin B, Nazar K, Costill DL, Stein E, Jansson E, Essen B, and Gollnick D. The nature of the training response; peripheral and central adaptations of one legged exercise. *Acta Physiol Scand* 96: 289–305, 1976.
- Savard GK, Areskog NH, and Saltin B. Maximal muscle activation is not limited by pulmonary ventilation in chronic hypoxia. *Acta Physiol Scand* 157: 187–190, 1996.
- Stenberg J, Eklblom B, and Messin R. Hemodynamic response to work at simulated altitude, 4,000 m. *J Appl Physiol* 21: 1589–1594, 1966.

Bengt Saltin  
 José A. L. Calbet  
 The Copenhagen Muscle Research Centre  
 Rigshospitalet  
 Copenhagen N, Denmark  
 Department of Physical Education  
 University of Las Palmas de Gran Canaria  
 Gran Canaria, Spain  
 e-mail: bengt.saltin@rh.hosp.dk

## Counterpoint: In health and in a normoxic environment, $\dot{V}O_{2\max}$ is not limited primarily by cardiac output and locomotor muscle blood flow

Let's begin this by being sure of the question we are addressing, because this topic is notorious for being easy to spin toward one's desired position by subtly changing the question. I would like to clear the deck of spin right from the

start. So I will stipulate that without blood flow,  $\dot{V}O_{2\max}$  would be zero: Saltin 1, Wagner 0. I will also stipulate that the venerable Fick Principle, taken at its naive simplest, would tend to support my opponent:  $\dot{V}O_2 = \dot{Q} \times [CaO_2 -$

$C_{vO_2}$ ], where  $\dot{Q}$  is cardiac output,  $C_{aO_2}$  is arterial, and  $C_{vO_2}$  mixed venous  $[O_2]$ .

I will even argue for him, comparing Lance Armstrong or equivalent with a sedentary normal subject each at their maximal exercise capacities,  $\dot{V}_{O_2}$  would be about twice as high in LA (~80 vs. ~40 ml·kg<sup>-1</sup>·min<sup>-1</sup>).  $C_{aO_2}$  in the absence of erythropoietin would be close to 20 ml/dl in each, maybe even lower in LA if he shows exercise-induced desaturation (1) plus the plasma volume expansion, common in trained athletes, that results in a reduced [Hb] (16).  $C_{vO_2}$  would be lower in LA, perhaps as low as 2 ml/dl (i.e., 90% extraction) (5), whereas in his unfit couch potato (CP) counterpart, maximal extraction might not exceed 70% (12), with  $C_{vO_2}$  therefore at 6 ml/dl. Thus in the Fick equation above, maximal  $[C_{aO_2} - C_{vO_2}]$  approximates 180 ml/l in LA and 140 ml/l in CP. This, in turn implies that LA's peak  $\dot{Q}$  must be 32 l/min, whereas CP's is only 20 l/min (assuming both weigh ~70 kg). For LA,  $\dot{Q}$  is 60% higher but  $[C_{aO_2} - C_{vO_2}]$  is only 30% higher. So Bengt would be justified in saying  $\dot{Q}$  is the primary determinant of  $\dot{V}_{O_2 \max}$  if the question is "what primarily explains the difference in  $\dot{V}_{O_2 \max}$  between CP and LA?  $\dot{Q}$  or  $[C_{aO_2} - C_{vO_2}]$ ?" Saltin 1.5, Wagner 0. (I will return to LA and CP later. Bengt, watch out.)

But, this is *not* the question that we are being asked to address. The question is: "Is cardiac output (or muscle blood flow) the primary determinant of  $\dot{V}_{O_2 \max}$  or not?" Stated in other words, if a normal subject is exercising at  $\dot{V}_{O_2 \max}$  and you were somehow able to augment any single part of the  $O_2$  transport and use chain, what effect would this have on  $\dot{V}_{O_2 \max}$ ? And, would cardiac output, as one part of that chain, have the largest effect, as Bengt will argue? I hope he will not try and argue  $\dot{Q}$  is the sole limiting factor, or I will blow him out of the water in rebuttal.

There is undeniable evidence that  $\dot{V}_{O_2 \max}$  can be acutely altered at will in normal humans by any one of a number of interventions (8, 10, 14, 17, 21), of which altering  $\dot{Q}$  is but one. Let's step down the  $O_2$  transport pathway, examining each step in turn.

Changing  $F_{I_{O_2}}$  changes  $\dot{V}_{O_2 \max}$  in the same direction (5, 6). Ventilation at  $\dot{V}_{O_2 \max}$  is very hard to alter in normal subjects, but published theoretical models demonstrate that maximal  $O_2$  transport and thus  $\dot{V}_{O_2 \max}$  would be affected by changes in ventilation (20).  $\dot{V}_A/\dot{Q}$  inequality (2), alveolar-capillary diffusion limitation (18), and (post) pulmonary shunts (2) can and do play a small but demonstrable role in reducing arterial oxygenation and thus  $\dot{V}_{O_2 \max}$ , as our own editor showed many years ago (9). Cardiac output (or muscle blood flow) clearly affects  $\dot{V}_{O_2 \max}$ , although direct interventions to test this have been done only in animals such as dogs, for example, by pericardiectomy (3), which allows a higher cardiac output and  $\dot{V}_{O_2 \max}$ . Changes in [Hb] (15) and in the  $P_{50}$  of Hb (4, 11) both alter convective  $O_2$  transport to the muscles and have been shown to affect  $\dot{V}_{O_2 \max}$  in controlled studies. Skeletal muscle  $O_2$  transport conductance (between capillaries and mitochondria), which relates closely to capillarity, has also been shown to play a significant role in setting  $\dot{V}_{O_2 \max}$  (13). Finally, maximal mitochondrial rate of  $O_2$  consumption has the power to affect  $\dot{V}_{O_2 \max}$  (7).

Although the above demonstrates, beyond argument even by Bengt, that  $\dot{Q}$  is by no means the only factor contributing to  $\dot{V}_{O_2 \max}$ , I have not yet provided the key arguments that must

address the core question of sensitivity of  $\dot{V}_{O_2 \max}$  to a given percent change in each of the above steps. Saltin still 1.5, Wagner still 0. Answering that question will put the nail in the  $\dot{Q}$ /Saltin coffin, as follows.

First, suppose maximal mitochondrial  $O_2$  consumption is *less* than maximal  $O_2$  available by transport from the air to the mitochondria. Further raising  $O_2$  transport by increasing cardiac output (or for that matter any of the other above  $O_2$  pathway steps) will have no effect on  $\dot{V}_{O_2 \max}$  because it is by definition  $O_2$  supply independent. Saltin 1.5, Wagner 1.0.

But suppose things are turned the other way around: maximal mitochondrial  $O_2$  use potential now *exceeds*  $O_2$  availability. Then, according to the evidence presented above, augmenting each and every step in  $O_2$  transport should have a positive effect on  $\dot{V}_{O_2 \max}$ , and it does. Suppose each component is augmented by 20% of its value, one at a time. Integrated physiological models incorporating all pathway steps (20) and Fig. 1 show that a 20% increase in  $F_{I_{O_2}}$  raises  $\dot{V}_{O_2 \max}$  by only 5.0%, due to the flat  $O_2$ -Hb dissociation curve in the normal range. Increasing ventilation 20% will also lead to a small (1.3%) increase, again because  $PO_2$  is on the flat part of the curve, and raising  $PO_2$  has little effect on  $C_{aO_2}$ . Increase lung diffusing capacity 20% in an athlete who has *mild* hypoxemia due to diffusion limitation and  $\dot{V}_{O_2 \max}$  will increase by 2.9%. Increasing diffusing capacity in a subject without diffusion limitation obviously cannot improve  $\dot{V}_{O_2 \max}$ . If skeletal muscle  $O_2$  diffusional conductance is increased by 20%,  $\dot{V}_{O_2 \max}$  will be 5.0% higher. Increase [Hb] by 20% and  $\dot{V}_{O_2 \max}$  increases by only 3.9%. Finally, increase  $\dot{Q}$  by 20%, and  $\dot{V}_{O_2 \max}$  increases by only 2.6%, *half that when muscle  $O_2$  conductance is raised equally*. Why? Because muscle  $O_2$  conductance has only one significant effect—to increase  $O_2$  flux from blood to cells. But raising  $\dot{Q}$  has opposing effects (19). First, it *increases* convective  $O_2$  transport by the circulation as predicted by both Bengt and the Fick principle. But the higher  $\dot{Q}$  simultaneously reduces transit time in both lung




Fig. 1. Calculated effects of individual changes in key  $O_2$  transport variables on  $\dot{V}_{O_2 \max}$ . Data reflect typical normal sea level values. Calculations use the model are described in Ref. 20. Note that all variables affect  $\dot{V}_{O_2}$ , and that  $\dot{Q}$  is by no means the most important factor.

and muscle capillaries and this *worsens* diffusion limitation, significantly opposing this convective gain.

This brings me back to LA and CP as promised. If LA did not have a superior muscle O<sub>2</sub> conductance to facilitate O<sub>2</sub> transport to cells, the 32 l/min  $\dot{Q}$  would simply limit O<sub>2</sub> extraction due to rapid red cell transit. *The only way LA can get to 80 ml/min  $\dot{V}_{O_{2\max}}$  is by having both an exceptional  $\dot{Q}$  and a matching, exceptional muscle capillary-to-mitochondrion O<sub>2</sub> transport system to permit almost full O<sub>2</sub> extraction from the rapidly flowing blood.* Thus, even if Bengt argues from the Fick Principle, as in my opening paragraph, the untold story is that muscle O<sub>2</sub> conductance must also be extraordinary, every bit as important as  $\dot{Q}$ , or O<sub>2</sub> extraction could not possible reach 90%. I rest my case, Bengt: Saltin 1.5, Wagner 10.

#### REFERENCES

- Dempsey JA, Hanson PG, and Henderson KS. Exercise-induced arterial hypoxemia in healthy human subjects at sea level. *J Physiol* 355: 161–175, 1984.
- Gledhill N, Froese AB, and Dempsey JA. Ventilation to perfusion distribution during exercise in health. In: *Muscular Exercise and the Lung*, edited by Dempsey JA and Reed CE. Madison: University of Wisconsin Press, 1977, p. 325–344.
- Hammond HK, White FC, Bhargava V, and Shabetai R. Heart size and maximal cardiac output are limited by the pericardium. *Am J Physiol Heart Circ Physiol* 263: H1675–H1681, 1992.
- Hogan MC, Bebout DE, and Wagner PD. Effect of increased Hb-O<sub>2</sub> affinity on  $\dot{V}_{O_{2\max}}$  at constant O<sub>2</sub> delivery in dog muscle in situ. *J Appl Physiol* 70: 2656–2662, 1991.
- Knight DR, Poole DC, Schaffartzik W, Guy HJ, Prediletto R, Hogan MC, and Wagner PD. Relationship between body and leg  $\dot{V}_{O_2}$  during maximal cycle ergometry. *J Appl Physiol* 73: 1114–1121, 1992.
- Knight DR, Schaffartzik W, Poole DC, Hogan MC, Bebout DE, and Wagner PD. Effects of hyperoxia on maximal leg O<sub>2</sub> supply and utilization in men. *J Appl Physiol* 75: 2586–2594, 1993.
- McAllister RM and Terjung RL. Acute inhibition of respiratory capacity of muscle reduces peak oxygen consumption. *Am J Physiol Cell Physiol* 259: C889–C896, 1990.
- Pirnay F, Lamy M, Dujardin J, Deroanne R, and Petit M. Analysis of femoral venous blood during maximum exercise. *J Appl Physiol* 33: 289–292, 1972.
- Powers SK, Lawler J, Dempsey J, Dodd JA, and Landry G. Effects of incomplete pulmonary gas exchange on  $\dot{V}_{O_{2\max}}$ . *J Appl Physiol* 66: 2491–2495, 1989.
- Pugh LGCE, Gill MB, Lahiri S, Milledge JS, Ward MP, and West JB. Muscular exercise at great altitudes. *J Appl Physiol* 19: 431–440, 1964.
- Richardson RS, Tagore K, Haseler L, Jordan M, and Wagner PD. Increased  $\dot{V}_{O_{2\max}}$  with a right shifted Hb-O<sub>2</sub> dissociation curve at a constant O<sub>2</sub> delivery in dog muscle in situ. *J Appl Physiol* 84: 995–1002, 1998.
- Roca J, Agustí AGN, Alonso A, Poole DC, Viegas C, Barberá JA, Rodríguez-Roisin R, Ferrer A, and Wagner PD. Effects of training on muscle O<sub>2</sub> transport at  $\dot{V}_{O_{2\max}}$ . *J Appl Physiol* 73: 1067–1076, 1992.
- Roca J, Hogan MC, Story D, Bebout DE, Haab P, Gonzalez R, Ueno O, and Wagner PD. Evidence for tissue diffusion limitation of  $\dot{V}_{O_{2\max}}$  in normal humans. *J Appl Physiol* 67: 291–299, 1989.
- Saltin B, Blomqvist CG, Mitchell JH, Johnson RL Jr, Wildenthal K, and Chapman CB. Response to exercise after bed rest and after training: a longitudinal study of adaptive changes in oxygen transport and body composition. *Circulation* 38, Suppl 7: 1–78, 1968.
- Schaffartzik W, Barton ED, Poole DC, Tsukimoto K, Hogan MC, Bebout DE, and Wagner PD. Effect of reduced hemoglobin concentration on leg oxygen uptake during maximal exercise in humans. *J Appl Physiol* 75: 491–498, 1993.
- Schumacher YO, Schmid A, Grathwohl D, Bultermann D, and Berg A. Hematological indices and iron status in athletes of various sports and performances. *Med Sci Sports Exerc* 34: 869–875, 2002.
- Spriet LL, Gledhill N, Froese AB, Wilkes DL, and Meyers EC. The effect of induced erythrocythemia on central circulation and oxygen transport during maximal exercise (Abstract). *Med Sci Sports Exerc* 12: 122, 1980.
- Torre-Bueno J, Wagner PD, Saltzman HA, Gale GE, and Moon RE. Diffusion limitation in normal humans during exercise at sea level and simulated altitude. *J Appl Physiol* 58: 989–995, 1985.
- Wagner PD. Algebraic analysis of the determinants of  $\dot{V}_{O_{2\max}}$ . *Respir Physiol* 93: 221–237, 1993.
- Wagner PD. A theoretical analysis of factors determining  $\dot{V}_{O_{2\max}}$  at sea level and altitude. *Respir Physiol* 106: 329–343, 1996.
- Welch HG. Hyperoxia and human performance: a brief review. *Med Sci Sports Exerc* 14: 253–262, 1982.

Peter D. Wagner  
University of California, San Diego  
La Jolla, California  
e-mail: pdwagner@ucsd.edu

#### REBUTTAL FROM DRS. SALTIN AND CALBET

The diffusional limitation theory is based primarily on one study (8) where an extraordinary elevation of leg  $\dot{V}_{O_{2\text{peak}}}$  (39%) and whole body (WB)  $\dot{V}_{O_{2\max}}$  (35%) was observed after 6 wk of training (Ref. 8, p. 1070), whereas maximal exercise intensity was only enhanced by 9%. Leg  $\dot{V}_{O_2}$  only accounted for 53–55% of WB  $\dot{V}_{O_2}$  at maximal exercise (before-after training), i.e., far below the normal 75–85% (5). These low leg peak  $\dot{V}_{O_2}$  values were likely caused by underestimation of peak leg blood flow (BF; which was only 5–6 l/min). Because during WB exercise, systemic a-v difference is never higher than leg a-v difference, peak cardiac output should have been >19 l/min before training and >23 l/min after training (+20%), leaving 9–10 l/min of BF for the rest of the body, which is too high a figure (5). Because DO<sub>2</sub> (oxygen conductance) is calculated as peak leg  $\dot{V}_{O_2}$ /mean capillary Po<sub>2</sub> (PmcO<sub>2</sub>) (10), it is likely that DO<sub>2</sub> was also underestimated (8).

Could a “couch potato” (CP) enhance his  $\dot{V}_{O_{2\max}}$  by increasing his cardiac output and BF? CP should be able to achieve an arm BF of ~2.5–3 l/min with an O<sub>2</sub> extraction a bit lower in the arms than the legs during maximal exercise (1–3, 7). This means that the  $\dot{V}_{O_{2\text{peak}}}$  of CP arms could reach 0.6–0.7 l/min. To perform maximal exercise with the four extremities, CP will need to increase his maximal cardiac output from 20 to 24 l/min. With the extra perfusion, CP could achieve a  $\dot{V}_{O_{2\max}}$  20% greater, even when assuming a lower muscle diffusing capacity in the arms than in the legs (2). CP could also increase his  $\dot{V}_{O_{2\max}}$  after blood transfusion or treatment with EPO. After this intervention, PmcO<sub>2</sub> will be similar or a bit higher (6), meaning that the increase of  $\dot{V}_{O_{2\max}}$  requires an increase of DO<sub>2</sub> after transfusion or EPO. If for a given PmcO<sub>2</sub>, DO<sub>2</sub> is enhanced when [Hb] is increased, it implies that  $\dot{V}_{O_{2\max}}$  is not limited by a structural resistance to diffusion in the skeletal muscle of healthy humans, i.e., what Roughton and Forster called membrane component of the oxygen conductance (9). Thus, for DO<sub>2</sub> to be the key limiting factor for  $\dot{V}_{O_{2\max}}$ , first the evidence that DO<sub>2</sub> actually represents the maximal attainable oxygen diffusing capacity in skeletal muscles should be provided. However, we agree that a diffusion limitation theoretically is a possibility but functionally it is a very minor player in healthy humans (4).

#### REFERENCES

- Ahlborg G and Jensen-Urstad M. Arm blood flow at rest and during arm exercise. *J Appl Physiol* 70: 928–933, 1991.
- Calbet JA, Holmberg HC, Rosdahl H, van Hall G, Jensen-Urstad M, and Saltin B. Why do arms extract less oxygen than legs during exercise? *Am J Physiol Regul Integr Comp Physiol* 289: R1448–R1458, 2005.

3. Calbet JA, Jensen-Urstad M, Van Hall G, Holmberg HC, Rosdahl H, and Saltin B. Maximal muscular vascular conductances during whole body upright exercise in humans. *J Physiol* 558: 319–331, 2004.
4. di Prampero PE and Ferretti G. Factors limiting maximal oxygen consumption in humans. *Respir Physiol* 80: 113–127, 1990.
5. Knight DR, Poole DC, Schaffartzik W, Guy HJ, Prediletto R, Hogan MC, and Wagner PD. Relationship between body and leg  $\dot{V}O_2$  during maximal cycle ergometry. *J Appl Physiol* 73: 1114–1121, 1992.
6. Marrades RM, Roca J, Campistol JM, Diaz O, Barbera JA, Torregrosa JV, Masclans JR, Cobos A, Rodriguez-Roisin R, and Wagner PD. Effects of erythropoietin on muscle O<sub>2</sub> transport during exercise in patients with chronic renal failure. *J Clin Invest* 97: 2092–2100, 1996.
7. Rasmussen B, Klausen K, Clausen JP, and Trap-Jensen J. Pulmonary ventilation, blood gases, and blood pH after training of the arms or the legs. *J Appl Physiol* 38: 250–256, 1975.
8. Roca J, Agusti AG, Alonso A, Poole DC, Viegas C, Barbera JA, Rodriguez-Roisin R, Ferrer A, and Wagner PD. Effects of training on muscle O<sub>2</sub> transport at  $\dot{V}O_{2\max}$ . *J Appl Physiol* 73: 1067–1076, 1992.
9. Roughton FJ and Forster RE. Relative importance of diffusion and chemical reaction rates in determining rate of exchange of gases in the human lung, with special reference to true diffusing capacity of pulmonary membrane and volume of blood in the lung capillaries. *J Appl Physiol* 11: 290–302, 1957.
10. Wagner PD. Algebraic analysis of the determinants of  $\dot{V}O_{2\max}$ . *Respir Physiol* 93: 221–237, 1993.

#### REBUTTAL FROM DR. WAGNER

My good friends Bengt and Jose have done a wonderful job of making my case and my rebuttal easy, because we clearly agree on several points. We agree that cardiac output/muscle blood flow is *one* determinant of maximal  $\dot{V}O_2$ . We agree that *one* major difference between an athlete and a couch potato is in maximal cardiac output. However, we surprisingly agree that other factors contribute substantially to maximal  $\dot{V}O_2$ . Bengt and Jose say this in *paragraph 2* referring to the role of

$CaO_2$ , which is *not* blood flow and restate this in their concluding paragraph, agreeing that lungs, heart, and blood are all important, just as I have argued. But they cannot use this to advance their own argument because the topic was *not* about O<sub>2</sub> delivery, it was about blood flow.

I must also remind my friends that the topic includes the word *primarily*. They provided *no* evidence that per unit of change in the responsible variable, blood flow is the primary factor, more important than any other conductances in the O<sub>2</sub> transport chain. They have failed to realize that for a high cardiac output to allow a high  $\dot{V}O_{2\max}$ , the diffusing capacities in both the lungs and muscles *must* be correspondingly high, or pulmonary O<sub>2</sub> loading and tissue unloading *must* be compromised, as pointed out many years ago by Piiper et al. (1, 2). They have assigned primary importance to one variable (flow) without assessing all other pertinent variables. How can you compare the roles of each variable when not all are addressed? Suppose you ask which is the fastest way to get from *point A* to *B*? By car, bicycle, or plane, and don't even study other alternatives such as by train or on foot. You simply cannot conclude that by train or on foot are not faster ways to get there. By looking at only part of the story, they have presented only part of the answer.

#### REFERENCES

1. Piiper J, Meyer M, and Scheid P. Dual role of diffusion in tissue gas exchange: blood-tissue equilibration and diffusion shunt. *Respir Physiol* 56: 131–144, 1984.
2. Piiper J and Scheid P. Model for capillary-alveolar equilibration with special reference to O<sub>2</sub> uptake in hypoxia. *Respir Physiol* 46: 193–208, 1981.